wwPDB X-ray Structure Validation Summary Report (i)

Mar 18, 2026 – 08:28 AM UTC

PDB ID	:	2BR8 / pdb_00002br8
Title	:	Crystal Structure of Acetylcholine-binding Protein (AChBP) from Aplysia californica in complex with an alpha-conotoxin PnIA variant
Authors	:	Celie, P.H.N.; Kasheverov, I.E.; Mordvintsev, D.Y.; Hogg, R.C.; van Nierop, P.; van Elk, R.; van Rossum-Fikkert, S.E.; Zhmak, M.N.; Bertrand, D.; Tsetlin, V.; Sixma, T.K.; Smit, A.B.
Deposited on	:	2005-05-03
Resolution	:	2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	217	2% 74% 19% • 6%
1	B	217	3% 79% 13% • 6%
1	C	217	5% 75% 18% • 6%
1	D	217	77% 16% • 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	217	
2	F	17	
2	G	17	
2	H	17	
2	I	17	
2	J	17	

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 9010 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SOLUBLE ACETYLCHOLINE RECEPTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	205	Total	C	N	O	S	0	0	0
			1636	1036	266	325	9			
1	B	205	Total	C	N	O	S	0	0	0
			1636	1036	266	325	9			
1	C	205	Total	C	N	O	S	0	0	0
			1636	1036	266	325	9			
1	D	205	Total	C	N	O	S	0	0	0
			1636	1036	266	325	9			
1	E	205	Total	C	N	O	S	0	0	0
			1636	1036	266	325	9			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	41	VAL	ALA	conflict	UNP Q8WSF8
A	136	VAL	ALA	conflict	UNP Q8WSF8
B	41	VAL	ALA	conflict	UNP Q8WSF8
B	136	VAL	ALA	conflict	UNP Q8WSF8
C	41	VAL	ALA	conflict	UNP Q8WSF8
C	136	VAL	ALA	conflict	UNP Q8WSF8
D	41	VAL	ALA	conflict	UNP Q8WSF8
D	136	VAL	ALA	conflict	UNP Q8WSF8
E	41	VAL	ALA	conflict	UNP Q8WSF8
E	136	VAL	ALA	conflict	UNP Q8WSF8

- Molecule 2 is a protein called ALPHA-CONOTOXIN PNIA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	17	Total	C	N	O	S	0	0	1
			114	70	20	20	4			
2	G	17	Total	C	N	O	S	0	0	1
			114	70	20	20	4			

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	17	Total	C	N	O	S	0	0	1
			114	70	20	20	4			
2	I	17	Total	C	N	O	S	0	0	1
			114	70	20	20	4			
2	J	17	Total	C	N	O	S	0	0	1
			114	70	20	20	4			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	10	LEU	ALA	engineered mutation	UNP P50984
F	14	LYS	ASP	engineered mutation	UNP P50984
G	10	LEU	ALA	engineered mutation	UNP P50984
G	14	LYS	ASP	engineered mutation	UNP P50984
H	10	LEU	ALA	engineered mutation	UNP P50984
H	14	LYS	ASP	engineered mutation	UNP P50984
I	10	LEU	ALA	engineered mutation	UNP P50984
I	14	LYS	ASP	engineered mutation	UNP P50984
J	10	LEU	ALA	engineered mutation	UNP P50984
J	14	LYS	ASP	engineered mutation	UNP P50984

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	E	1	Total O S 5 4 1	0	0
3	F	1	Total O S 5 4 1	0	0
3	G	1	Total O S 5 4 1	0	0
3	I	1	Total O S 5 4 1	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	35	Total O 35 35	0	0
4	B	38	Total O 38 38	0	0
4	C	71	Total O 71 71	0	0
4	D	51	Total O 51 51	0	0
4	E	26	Total O 26 26	0	0
4	G	4	Total O 4 4	0	0
4	H	5	Total O 5 5	0	0
4	I	4	Total O 4 4	0	0
4	J	1	Total O 1 1	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: SOLUBLE ACETYLCHOLINE RECEPTOR

Chain A:




• Molecule 1: SOLUBLE ACETYLCHOLINE RECEPTOR

Chain B:




• Molecule 1: SOLUBLE ACETYLCHOLINE RECEPTOR

Chain C:



• Molecule 1: SOLUBLE ACETYLCHOLINE RECEPTOR

Chain D:






• Molecule 1: SOLUBLE ACETYLCHOLINE RECEPTOR

Chain E:






• Molecule 2: ALPHA-CONOTOXIN PNIA

Chain F:






• Molecule 2: ALPHA-CONOTOXIN PNIA

Chain G:





• Molecule 2: ALPHA-CONOTOXIN PNIA

Chain H:





• Molecule 2: ALPHA-CONOTOXIN PNIA

Chain I:






• Molecule 2: ALPHA-CONOTOXIN PNIA

Chain J:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	67.47Å 80.08Å 134.04Å 90.00° 93.20° 90.00°	Depositor
Resolution (Å)	19.80 – 2.40 19.80 – 2.40	Depositor EDS
% Data completeness (in resolution range)	97.8 (19.80-2.40) 97.6 (19.80-2.40)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.19 (at 2.41Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.198 , 0.250 (Not available) , 0.249	Depositor DCC
R_{free} test set	2783 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å^2)	36.8	Xtriage
Anisotropy	0.221	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 29.0	EDS
L-test for twinning[2]	$< \|L\| > = 0.47$, $< L^2 > = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.92	EDS
Total number of atoms	9010	wwPDB-VP
Average B, all atoms (Å^2)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|>5$	RMSZ	#$\|Z\|>5$
1	A	0.54	0/1676	0.88	0/2287
1	B	0.62	0/1676	0.92	3/2287 (0.1%)
1	C	0.72	0/1676	1.00	4/2287 (0.2%)
1	D	0.63	0/1676	0.94	3/2287 (0.1%)
1	E	0.58	0/1676	0.93	2/2287 (0.1%)
2	F	0.55	0/116	1.09	0/158
2	G	0.68	0/116	1.01	1/158 (0.6%)
2	H	0.75	0/116	1.00	1/158 (0.6%)
2	I	0.55	0/116	0.97	0/158
2	J	0.55	0/116	1.05	0/158
All	All	0.62	0/8960	0.94	14/12225 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	2
1	D	0	1
All	All	0	4

There are no bond length outliers.

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	45	THR	N-CA-C	-6.69	100.06	110.17
1	E	45	THR	N-CA-C	-6.29	101.16	110.46
1	C	3	ASN	N-CA-C	-6.28	105.62	113.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	C	45	THR	N-CA-C	-6.05	101.04	110.42
1	D	45	THR	CA-C-N	-6.04	114.86	123.20

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	16	PRO	Peptide
1	C	16	PRO	Peptide
1	C	188	CYS	Peptide
1	D	15	SER	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1636	0	1572	26	0
1	B	1636	0	1572	27	0
1	C	1636	0	1572	23	0
1	D	1636	0	1572	22	0
1	E	1636	0	1572	14	0
2	F	114	0	108	5	0
2	G	114	0	108	4	0
2	H	114	0	108	1	0
2	I	114	0	108	4	0
2	J	114	0	108	2	0
3	C	5	0	0	0	0
3	E	5	0	0	0	0
3	F	5	0	0	0	0
3	G	5	0	0	0	0
3	I	5	0	0	0	0
4	A	35	0	0	1	0
4	B	38	0	0	4	0
4	C	71	0	0	4	0
4	D	51	0	0	4	0
4	E	26	0	0	0	0
4	G	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	H	5	0	0	0	0
4	I	4	0	0	0	0
4	J	1	0	0	0	0
All	All	9010	0	8400	107	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 107 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:114:MET:HE1	1:E:116:ILE:HD11	1.49	0.93
1:B:95:ARG:HG2	1:B:95:ARG:HH11	1.38	0.87
1:B:50:LEU:HB3	4:B:2008:HOH:O	1.73	0.86
1:B:154:LEU:O	1:B:181:ARG:HD2	1.76	0.85
1:D:16:PRO:HD3	4:D:2004:HOH:O	1.77	0.82

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	203/217 (94%)	198 (98%)	3 (2%)	2 (1%)	12 20
1	B	203/217 (94%)	192 (95%)	10 (5%)	1 (0%)	24 37
1	C	203/217 (94%)	192 (95%)	9 (4%)	2 (1%)	12 20
1	D	203/217 (94%)	194 (96%)	6 (3%)	3 (2%)	8 12
1	E	203/217 (94%)	195 (96%)	7 (3%)	1 (0%)	24 37
2	F	15/17 (88%)	14 (93%)	1 (7%)	0	100 100
2	G	15/17 (88%)	15 (100%)	0	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	H	15/17 (88%)	15 (100%)	0	0	100	100
2	I	15/17 (88%)	14 (93%)	1 (7%)	0	100	100
2	J	15/17 (88%)	15 (100%)	0	0	100	100
All	All	1090/1170 (93%)	1044 (96%)	37 (3%)	9 (1%)	16	25

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2	ALA
1	C	15	SER
1	D	15	SER
1	E	15	SER
1	B	16	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/197 (95%)	176 (94%)	12 (6%)	16	28
1	B	188/197 (95%)	178 (95%)	10 (5%)	20	36
1	C	188/197 (95%)	178 (95%)	10 (5%)	20	36
1	D	188/197 (95%)	176 (94%)	12 (6%)	16	28
1	E	188/197 (95%)	180 (96%)	8 (4%)	26	44
2	F	14/14 (100%)	14 (100%)	0	100	100
2	G	14/14 (100%)	13 (93%)	1 (7%)	13	23
2	H	14/14 (100%)	13 (93%)	1 (7%)	13	23
2	I	14/14 (100%)	14 (100%)	0	100	100
2	J	14/14 (100%)	14 (100%)	0	100	100
All	All	1010/1055 (96%)	956 (95%)	54 (5%)	20	36

5 of 54 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	169	SER
1	D	68	ASN
1	E	178	THR
1	C	180	THR
1	D	5	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	184	GLN
1	E	72	ASN
1	D	55	GLN
1	E	184	GLN
1	E	46	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	SO4	G	1018	-	4,4,4	0.25	0	6,6,6	0.14	0
3	SO4	E	1206	-	4,4,4	0.22	0	6,6,6	0.09	0
3	SO4	I	1018	-	4,4,4	0.22	0	6,6,6	0.23	0
3	SO4	F	1018	-	4,4,4	0.22	0	6,6,6	0.10	0
3	SO4	C	1206	-	4,4,4	0.28	0	6,6,6	0.27	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	J	1
2	H	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	J	16:CYS	C	17:NH2	N	1.65
1	H	16:CYS	C	17:NH2	N	1.04

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	205/217 (94%)	0.09	5 (2%) 59 55	18, 25, 41, 52	0
1	B	205/217 (94%)	-0.05	7 (3%) 48 44	11, 19, 48, 56	0
1	C	205/217 (94%)	0.11	10 (4%) 35 31	10, 21, 47, 60	0
1	D	205/217 (94%)	-0.39	0 100 100	7, 15, 24, 38	0
1	E	205/217 (94%)	-0.14	3 (1%) 72 68	10, 17, 37, 44	0
2	F	16/17 (94%)	-0.06	0 100 100	21, 23, 27, 27	0
2	G	16/17 (94%)	-0.14	0 100 100	19, 22, 24, 25	0
2	H	16/17 (94%)	-0.20	0 100 100	13, 17, 19, 20	0
2	I	16/17 (94%)	-0.02	0 100 100	23, 25, 27, 27	0
2	J	16/17 (94%)	0.38	0 100 100	31, 33, 35, 35	0
All	All	1105/1170 (94%)	-0.07	25 (2%) 61 57	7, 20, 40, 60	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	16	PRO	5.7
1	C	15	SER	4.3
1	A	2	ALA	3.9
1	B	68	ASN	3.6
1	B	158	THR	3.4

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	SO4	E	1206	5/5	0.71	0.20	90,91,91,91	0
3	SO4	F	1018	5/5	0.85	0.16	76,76,76,77	0
3	SO4	G	1018	5/5	0.97	0.13	38,38,40,40	0
3	SO4	I	1018	5/5	0.97	0.08	27,27,28,28	0
3	SO4	C	1206	5/5	0.98	0.07	18,19,20,20	0

6.5 Other polymers (i)

There are no such residues in this entry.