Full wwPDB X-ray Structure Validation Report (i)

Aug 24, 2026 – 07:25 pm BST

PDB ID	:	9ERR / pdb_00009err
Title	:	Hydrogenase-2 Ni-SCO state
Authors	:	Carr, S.B.; Li, W.; Wong, K.l.; Ash, P.A.; Vincent, K.A.
Deposited on	:	2024-03-25
Resolution	:	1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	S	298	2% 85% 5% 10%
1	T	298	3% 87% • 10%
2	L	567	% 92% 5% •
2	M	567	2% 92% 5% •

2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 14020 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hydrogenase-2 small chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	S	269	2083	1318	368	384	13	0	5	0
1	T	269	2083	1320	366	384	13	0	6	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
S	294	HIS	-	expression tag	UNP P69741
S	295	HIS	-	expression tag	UNP P69741
S	296	HIS	-	expression tag	UNP P69741
S	297	HIS	-	expression tag	UNP P69741
S	298	HIS	-	expression tag	UNP P69741
S	299	HIS	-	expression tag	UNP P69741
T	294	HIS	-	expression tag	UNP P69741
T	295	HIS	-	expression tag	UNP P69741
T	296	HIS	-	expression tag	UNP P69741
T	297	HIS	-	expression tag	UNP P69741
T	298	HIS	-	expression tag	UNP P69741
T	299	HIS	-	expression tag	UNP P69741

- Molecule 2 is a protein called Hydrogenase-2 large chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	551	4325	2751	741	815	18	0	6	0
2	M	551	4298	2735	739	806	18	0	2	0

- Molecule 3 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe_4S_4).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Fe	S		
3	S	1	8	4	4	0	0
3	S	1	8	4	4	0	0
3	T	1	8	4	4	0	0
3	T	1	8	4	4	0	0

- Molecule 4 is FE3-S4 CLUSTER (CCD ID: F3S) (formula: Fe_3S_4).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Fe	S		
4	S	1	7	3	4	0	0
			Total	Fe	S		
4	T	1	7	3	4	0	0

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	S	1	6	3	3	0	0
			Total	C	O		
5	T	1	6	3	3	0	0

- Molecule 6 is CARBONMONOXIDE-(DICYANO) IRON (CCD ID: FCO) (formula: C_3FeN_2O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
6	L	1	7	3	1	2	1	0	0
6	M	1	7	3	1	2	1	0	0

- Molecule 7 is NICKEL (II) ION (CCD ID: NI) (formula: Ni) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ni		
7	L	1	1	1	0	0
7	M	1	1	1	0	0

- Molecule 8 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
8	L	2	2	2	0	0
8	M	3	3	3	0	0

- Molecule 9 is CARBON MONOXIDE (CCD ID: CMO) (formula: CO) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
9	L	1	2	1	1	0	0
			Total	C	O		
9	M	1	2	1	1	0	0

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
10	S	179	179	179	0	0
			Total	O		
10	L	425	425	425	0	0
			Total	O		
10	T	167	167	167	0	0
			Total	O		
10	M	377	377	377	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Hydrogenase-2 small chain

Chain S:

• Molecule 1: Hydrogenase-2 small chain

Chain T:

• Molecule 2: Hydrogenase-2 large chain

Chain L:





• Molecule 2: Hydrogenase-2 large chain

Chain M:






D544
P545
C546
C549
H552
VAL
VAL
ASP
ALA
ASP
GLY
ASN
GLU
VAL
VAL
SER
VAL
LYS
VAL
LEU

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	100.19Å 99.75Å 168.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	64.50 – 1.40 64.50 – 1.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (64.50-1.40) 99.9 (64.50-1.40)	Depositor EDS
R_{merge}	0.26	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.31 (at 1.40Å)	Xtriage
Refinement program	REFMAC 5.8.0419	Depositor
R, R_{free}	0.161 , 0.181 0.162 , 0.181	Depositor DCC
R_{free} test set	16420 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å^2)	11.7	Xtriage
Anisotropy	0.046	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 34.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	0.005 for k,h,-l	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	14020	wwPDB-VP
Average B, all atoms (Å^2)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.58% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CMO, NI, FCO, SF4, MG, F3S

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	S	0.69	0/2147	1.00	8/2923 (0.3%)
1	T	0.68	0/2156	1.00	3/2935 (0.1%)
2	L	0.68	0/4444	0.98	5/6057 (0.1%)
2	M	0.68	0/4411	1.00	4/6013 (0.1%)
All	All	0.68	0/13158	0.99	20/17928 (0.1%)

There are no bond length outliers.

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	M	382	ASN	N-CA-CB	-7.94	98.50	110.01
2	L	99	HIS	CA-CB-CG	-7.86	105.94	113.80
2	L	382	ASN	N-CA-CB	-7.28	99.42	110.12
2	M	99	HIS	CA-CB-CG	-6.75	107.05	113.80
1	S	212	ASP	CA-CB-CG	6.62	119.22	112.60
2	L	382	ASN	OD1-CG-ND2	5.99	128.59	122.60
1	S	153	GLY	CA-C-N	-5.35	116.97	122.85
1	S	153	GLY	C-N-CA	-5.35	116.97	122.85
1	S	217	GLU	CB-CG-CD	5.34	121.67	112.60
1	S	66	HIS	CA-CB-CG	-5.29	108.51	113.80
1	T	81	ASP	CA-CB-CG	5.28	117.88	112.60
2	M	171	TYR	CB-CA-C	5.28	117.72	111.22
1	S	33	THR	CA-C-N	-5.12	117.61	122.37
1	S	33	THR	C-N-CA	-5.12	117.61	122.37
1	T	33	THR	CA-C-N	-5.11	117.61	122.37
1	T	33	THR	C-N-CA	-5.11	117.61	122.37
2	L	299	PHE	CA-CB-CG	5.10	118.90	113.80
2	M	544	ASP	CA-CB-CG	5.06	117.66	112.60
1	S	81	ASP	CA-CB-CG	5.03	117.63	112.60
2	L	544	ASP	CA-CB-CG	5.02	117.62	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	S	2083	0	2005	8	0
1	T	2083	0	2013	7	0
2	L	4325	0	4261	19	0
2	M	4298	0	4240	12	0
3	S	16	0	0	0	0
3	T	16	0	0	0	0
4	S	7	0	0	0	0
4	T	7	0	0	0	0
5	S	6	0	8	0	0
5	T	6	0	8	0	0
6	L	7	0	0	0	0
6	M	7	0	0	0	0
7	L	1	0	0	0	0
7	M	1	0	0	0	0
8	L	2	0	0	0	0
8	M	3	0	0	0	0
9	L	2	0	0	1	0
9	M	2	0	0	1	0
10	L	425	0	0	6	0
10	M	377	0	0	0	0
10	S	179	0	0	4	0
10	T	167	0	0	2	0
All	All	14020	0	12535	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:S:213:GLU:OE2	10:S:401:HOH:O	1.79	0.99

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:T:109:GLU:HG2	10:T:556:HOH:O	1.78	0.82
2:L:196[A]:GLU:OE2	10:L:701:HOH:O	2.01	0.77
2:M:273:LEU:HD13	2:M:413:LEU:HD21	1.77	0.67
2:L:374[B]:THR:HG21	2:L:516:GLY:HA2	1.74	0.67
1:S:106:LYS:HD3	10:S:562:HOH:O	1.95	0.66
2:L:305[A]:GLU:CD	2:L:373:LYS:HZ2	2.08	0.62
2:L:481:MET:HE3	10:L:705:HOH:O	2.00	0.62
2:M:313:ARG:HD3	2:M:322:TYR:CE2	2.41	0.55
2:L:305[A]:GLU:CD	2:L:373:LYS:NZ	2.65	0.54
1:S:70:GLU:HG2	10:S:471:HOH:O	2.08	0.54
2:L:391:ARG:O	2:L:395[A]:GLN:HG3	2.09	0.52
2:L:303:TYR:CE2	2:L:305[B]:GLU:CG	2.95	0.50
1:S:266:HIS:NE2	10:S:402:HOH:O	2.34	0.50
2:L:374[B]:THR:CG2	2:L:516:GLY:HA2	2.42	0.49
2:M:159:VAL:HG22	2:M:164:LEU:HD11	1.95	0.48
2:M:546:CYS:CB	9:M:604:CMO:C	2.90	0.48
1:T:109:GLU:CG	10:T:556:HOH:O	2.51	0.47
2:L:379:PRO:HA	10:L:1051:HOH:O	2.15	0.47
2:M:273:LEU:CD1	2:M:413:LEU:HD21	2.43	0.46
2:M:500:VAL:CG1	2:M:501:PRO:HD2	2.45	0.46
2:M:468:GLU:HA	2:M:488:ILE:O	2.17	0.45
2:L:337:LYS:HD2	10:L:1103:HOH:O	2.17	0.44
2:L:479:ARG:CZ	9:L:604:CMO:O	2.65	0.44
1:T:23:THR:HB	2:M:63:VAL:HG23	2.00	0.44
2:L:356:TRP:CD1	2:L:356:TRP:C	2.96	0.43
2:M:320:ASP:OD1	2:M:322:TYR:HB3	2.18	0.43
1:S:23:THR:HB	2:L:63:VAL:HG23	2.00	0.43
2:M:115:ASP:HB3	2:M:538:ARG:HG2	2.00	0.43
2:M:500:VAL:HG11	2:M:549:CYS:HB3	2.01	0.42
2:L:159:VAL:HG22	2:L:164:LEU:HD11	2.01	0.42
1:S:202:ASP:OD2	1:T:202:ASP:OD2	2.37	0.42
2:L:82:ILE:HD11	2:L:461:PRO:HB3	2.02	0.42
1:T:84:ILE:HD12	1:T:84:ILE:N	2.35	0.42
1:S:228:GLY:N	1:S:229:PRO:CD	2.83	0.42
1:T:247:TRP:CZ2	1:T:249:VAL:HB	2.55	0.41
2:L:156:LYS:HE2	10:L:849:HOH:O	2.20	0.41
2:L:500:VAL:HG11	2:L:549:CYS:HB3	2.02	0.41
1:S:196:GLU:HA	1:T:196:GLU:HA	2.03	0.41
2:L:178:LYS:HE2	10:L:720:HOH:O	2.20	0.41
2:M:133:MET:HE1	2:M:183:VAL:CG2	2.51	0.41
2:L:500:VAL:CG1	2:L:501:PRO:HD2	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	S	272/298 (91%)	262 (96%)	10 (4%)	0	100	100
1	T	273/298 (92%)	263 (96%)	10 (4%)	0	100	100
2	L	555/567 (98%)	532 (96%)	22 (4%)	1 (0%)	43	19
2	M	551/567 (97%)	528 (96%)	22 (4%)	1 (0%)	43	19
All	All	1651/1730 (95%)	1585 (96%)	64 (4%)	2 (0%)	48	21

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	M	211	LYS
2	L	211	LYS

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	S	219/239 (92%)	218 (100%)	1 (0%)	81	61
1	T	220/239 (92%)	218 (99%)	2 (1%)	70	44
2	L	471/479 (98%)	468 (99%)	3 (1%)	78	56
2	M	467/479 (98%)	463 (99%)	4 (1%)	70	44
All	All	1377/1436 (96%)	1367 (99%)	10 (1%)	76	52

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	S	251	ILE
2	L	102	HIS
2	L	150	LYS
2	L	512	ASN
1	T	65	LYS
1	T	251	ILE
2	M	2	SER
2	M	102	HIS
2	M	181	PRO
2	M	312	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	S	10	GLN
1	S	20	GLN
1	S	60	GLN
1	S	274	ASN
2	L	3	GLN
2	L	47	ASN
2	L	87	ASN
2	L	436	GLN
2	L	449	ASN
1	T	60	GLN
2	M	449	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 19 ligands modelled in this entry, 7 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	SF4	T	301	1	0,12,12	-	-	-		
3	SF4	T	303	1	0,12,12	-	-	-		
3	SF4	S	303	1	0,12,12	-	-	-		
3	SF4	S	301	1	0,12,12	-	-	-		
6	FCO	M	601	2	1,6,6	2.15	1 (100%)	-		
5	GOL	S	304	-	5,5,5	0.27	0	5,5,5	0.71	0
4	F3S	T	302	1	0,9,9	-	-	-		
4	F3S	S	302	1	0,9,9	-	-	-		
6	FCO	L	601	2	1,6,6	2.84	1 (100%)	-		
9	CMO	L	604	-	0,1,1	-	-	-		
5	GOL	T	304	-	5,5,5	0.31	0	5,5,5	0.55	0
9	CMO	M	604	-	0,1,1	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SF4	T	301	1	-	-	0/6/5/5
3	SF4	T	303	1	-	-	0/6/5/5
3	SF4	S	303	1	-	-	0/6/5/5
3	SF4	S	301	1	-	-	0/6/5/5
5	GOL	S	304	-	-	0/4/4/4	-
4	F3S	T	302	1	-	-	0/3/3/3
4	F3S	S	302	1	-	-	0/3/3/3
5	GOL	T	304	-	-	0/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	L	601	FCO	O3-C3	-2.84	1.12	1.16
6	M	601	FCO	O3-C3	-2.15	1.13	1.16

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	L	604	CMO	1	0
9	M	604	CMO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand FCO M 601



Bond lengths



Bond angles



Torsions



Rings

Ligand FCO L 601



Ligand CMO L 604



Ligand CMO M 604	
Bond lengths	Bond angles
 Torsions	Rings

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	S	269/298 (90%)	-0.13	7 (2%)	57	60	5, 13, 27, 47	5 (1%)
1	T	269/298 (90%)	0.02	8 (2%)	52	55	7, 14, 29, 49	6 (2%)
2	L	551/567 (97%)	-0.35	4 (0%)	84	87	6, 11, 23, 33	6 (1%)
2	M	551/567 (97%)	-0.12	11 (1%)	65	68	7, 13, 27, 43	2 (0%)
All	All	1640/1730 (94%)	-0.18	30 (1%)	67	71	5, 13, 26, 49	19 (1%)

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	M	2	SER	4.7
1	T	9	PRO	4.2
1	S	9	PRO	3.6
2	M	27	VAL	3.5
1	T	8	ASN	3.2
2	M	166	ILE	3.2
2	M	137	VAL	3.2
2	L	136	GLY	3.1
2	L	137	VAL	2.9
2	M	413	LEU	2.8
1	S	8	ASN	2.7
2	M	167	PHE	2.7
1	T	145	GLY	2.6
1	S	276	GLU	2.6
1	T	86	LEU	2.6
2	M	160	ALA	2.5
1	S	171	TYR	2.5
1	S	66	HIS	2.4
1	S	275	VAL	2.3
2	M	25	ASN	2.3
1	T	89	ASN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	S	172	GLY	2.3
1	T	66	HIS	2.3
2	M	515	VAL	2.2
1	T	276	GLU	2.2
2	M	469	PHE	2.2
2	M	172	TRP	2.1
1	T	96	ALA	2.1
2	L	515	VAL	2.0
2	L	270	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
8	MG	M	606	1/1	0.87	0.20	43,43,43,43	0
9	CMO	M	604	2/2	0.87	0.23	20,20,20,24	0
8	MG	L	605	1/1	0.91	0.11	30,30,30,30	0
9	CMO	L	604	2/2	0.93	0.22	18,18,18,20	0
5	GOL	T	304	6/6	0.95	0.09	17,22,22,24	0
8	MG	M	605	1/1	0.96	0.07	18,18,18,18	0
5	GOL	S	304	6/6	0.96	0.07	18,20,22,22	0
3	SF4	S	303	8/8	1.00	0.02	8,9,9,9	0
6	FCO	L	601	7/7	1.00	0.03	7,8,8,9	0
6	FCO	M	601	7/7	1.00	0.03	8,8,10,11	0
7	NI	L	602	1/1	1.00	0.02	10,10,10,10	0
7	NI	M	602	1/1	1.00	0.03	11,11,11,11	0
8	MG	L	603	1/1	1.00	0.05	4,4,4,4	0
3	SF4	T	301	8/8	1.00	0.02	10,11,11,11	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
8	MG	M	603	1/1	1.00	0.06	6,6,6,6	0
3	SF4	T	303	8/8	1.00	0.02	9,10,11,11	0
4	F3S	S	302	7/7	1.00	0.02	9,9,9,10	0
4	F3S	T	302	7/7	1.00	0.02	9,9,10,10	0
3	SF4	S	301	8/8	1.00	0.02	10,11,11,11	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around CMO M 604:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around CMO L 604:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FCO L 601:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FCO M 601:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around NI L 602:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around NI M 602:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers (i)

There are no such residues in this entry.