wwPDB X-ray Structure Validation Summary Report (i)

Aug 17, 2026 – 11:24 pm BST

PDB ID	:	5JSU / pdb_00005jsu
Title	:	The 3D structure of the U489C variant of [NiFeSe] hydrogenase from Desulfovibrio vulgaris Hildenborough in the oxidized state at 1.40 Angstrom resolution
Authors	:	Marques, M.C.; Pereira, I.A.C.; Matias, P.M.
Deposited on	:	2016-05-09
Resolution	:	1.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	317	2% 83% 7% 11%
2	B	508	91% · 5%

2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 12971 atoms, of which 6103 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Periplasmic [NiFeSe] hydrogenase, small subunit.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	283	4440	1428	2203	371	418	20	1	24	0


- Molecule 2 is a protein called Periplasmic [NiFeSe] hydrogenase, large subunit, selenocysteine-containing.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	482	7775	2468	3900	674	711	22	0	23	0

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	4	TRP	-	expression tag	UNP Q72AS3
B	5	SER	-	expression tag	UNP Q72AS3
B	6	HIS	-	expression tag	UNP Q72AS3
B	7	PRO	-	expression tag	UNP Q72AS3
B	8	GLN	-	expression tag	UNP Q72AS3
B	9	PHE	-	expression tag	UNP Q72AS3
B	10	GLU	-	expression tag	UNP Q72AS3
B	11	LYS	-	expression tag	UNP Q72AS3
B	489	CSD	SEC	engineered mutation	UNP Q72AS3

- Molecule 3 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe_4S_4).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Fe S 8 4 4	0	0
3	A	1	Total Fe S 8 4 4	0	0
3	A	1	Total Fe S 8 4 4	0	1

- Molecule 4 is oxygen-damaged SF4 (CCD ID: 6ML) (formula: $Fe_4O_2S_4$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Fe O S 10 4 2 4	0	1

- Molecule 5 is CARBONMONOXIDE-(DICYANO) IRON (CCD ID: FCO) (formula: C_3FeN_2O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	Fe	N	O	0	0
			7	3	1	2	1		

- Molecule 6 is HYDROSULFURIC ACID (CCD ID: H2S) (formula: H_2S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	S	0	1
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total S 1 1	0	0

- Molecule 7 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total Ni 1 1	0	1

- Molecule 8 is FE (II) ION (CCD ID: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	B	1	Total Fe 1 1	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	280	Total O 280 280	0	0
9	B	431	Total O 431 431	0	1


3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Periplasmic [NiFeSe] hydrogenase, small subunit



• Molecule 2: Periplasmic [NiFeSe] hydrogenase, large subunit, selenocysteine-containing



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	106.20Å 62.66Å 109.84Å 90.00° 104.97° 90.00°	Depositor
Resolution (Å)	45.36 – 1.40 45.36 – 1.40	Depositor EDS
% Data completeness (in resolution range)	99.5 (45.36-1.40) 99.4 (45.36-1.40)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.11 (at 1.40Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.110 , 0.141 0.121 , 0.150	Depositor DCC
R_{free} test set	6833 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å^2)	9.6	Xtriage
Anisotropy	0.063	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.40 , 45.3	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.98	EDS
Total number of atoms	12971	wwPDB-VP
Average B, all atoms (Å^2)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.15% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CSX, FCO, FE2, NI, SF4, 6ML, CSD, H2S

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.36	0/2360	0.54	0/3204
2	B	0.33	0/3996	0.57	0/5394
All	All	0.34	0/6356	0.56	0/8598

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2237	2203	2134	15	2
2	B	3875	3900	3848	10	0
3	A	24	0	0	0	0
4	A	10	0	0	0	0
5	B	7	0	0	0	0
6	B	2	0	0	0	0
7	B	1	0	0	0	0
8	B	1	0	0	0	0
9	A	280	0	0	6	1
9	B	431	0	0	1	0
All	All	6868	6103	5982	24	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 24 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2:THR:O	9:A:401:HOH:O	2.04	0.74
1:A:213:TYR:OH	1:A:261[B]:GLN:OE1	2.06	0.74
1:A:218:ASP:OD2	9:A:403:HOH:O	2.12	0.66
1:A:21[B]:CYS:SG	1:A:118:VAL:HG12	2.39	0.61
2:B:453:ASP:OD1	2:B:455[B]:MET:HG2	2.04	0.58

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:178[B]:LYS:HZ1	1:A:209:GLU:OE1[2_658]	1.56	0.04
1:A:1:GLY:N	9:A:403:HOH:O[2_658]	2.17	0.03

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	304/317 (96%)	295 (97%)	8 (3%)	1 (0%)	36 16
2	B	500/508 (98%)	491 (98%)	9 (2%)	0	100 100
All	All	804/825 (98%)	786 (98%)	17 (2%)	1 (0%)	48 21

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	263	ALA

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	246/253 (97%)	246 (100%)	0	100 100
2	B	413/412 (100%)	406 (98%)	7 (2%)	53 21
All	All	659/665 (99%)	652 (99%)	7 (1%)	73 37

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	152[A]	LYS
2	B	152[B]	LYS
2	B	403	VAL
2	B	224	VAL
2	B	80	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	ASN
1	A	180	ASN
2	B	132	GLN
2	B	255	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	CSD	B	489	6,7,2	3,7,8	1.00	0	1,8,10	2.04	1 (100%)
2	CSX	B	75[C]	2	3,6,7	1.20	0	1,6,8	1.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CSD	B	489	6,7,2	-	1/2/6/8	-
2	CSX	B	75[C]	2	-	0/1/5/7	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	489	CSD	OD1-SG-CB	2.04	109.42	105.54

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	489	CSD	N-CA-CB-SG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 9 ligands modelled in this entry, 2 are modelled with single atom and 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	FCO	B	601	6,2	1,6,6	2.50	1 (100%)	-		
3	SF4	A	303[A]	1	0,12,12	-	-	-		
3	SF4	A	301	1	0,12,12	-	-	-		
3	SF4	A	302	1	0,12,12	-	-	-		
4	6ML	A	304[B]	1	0,12,12	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SF4	A	301	1	-	-	0/6/5/5
3	SF4	A	303[A]	1	-	-	0/6/5/5
4	6ML	A	304[B]	1	-	-	0/2/3/3
3	SF4	A	302	1	-	-	0/6/5/5

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	601	FCO	O3-C3	-2.50	1.12	1.16

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	283/317 (89%)	-0.91	5 (1%) 67 71	4, 12, 25, 56	18 (6%)
2	B	480/508 (94%)	-1.01	0 100 100	5, 12, 24, 36	19 (3%)
All	All	763/825 (92%)	-0.97	5 (0%) 84 87	4, 12, 24, 56	37 (4%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	GLY	5.6
1	A	2	THR	4.7
1	A	3	LEU	3.3
1	A	4	THR	3.0
1	A	5	GLY	2.6

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	CSX	B	75[C]	7/8	-	-	9,10,17,17	11
2	CSD	B	489	8/9	0.99	0.03	6,9,14,14	3

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
6	H2S	B	602[A]	1/1	0.99	0.03	12,12,12,12	1
7	NI	B	603[D]	1/1	0.99	0.06	11,11,11,11	1
3	SF4	A	303[A]	8/8	1.00	0.01	6,7,7,8	8
4	6ML	A	304[B]	10/10	1.00	0.03	6,7,8,8	10
5	FCO	B	601	7/7	1.00	0.03	7,8,10,12	0
3	SF4	A	301	8/8	1.00	0.01	9,9,10,10	0
6	H2S	B	605	1/1	1.00	0.02	7,7,7,7	0
3	SF4	A	302	8/8	1.00	0.01	7,7,7,7	0
8	FE2	B	604	1/1	1.00	0.01	8,8,8,8	0

6.5 Other polymers (i)

There are no such residues in this entry.